

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

TTF-1 Antibody / NKX2.1, Clone: [NX2.1/690], Mouse, Monoclonal NSJ-V2894-100UG

Artikelname	TTF-1 Antibody / NKX2.1, Clone: [NX2.1/690], Mouse, Monoclonal
Artikelnummer	NSJ-V2894-100UG
Hersteller Artikelnummer	V2894-100UG
Alternativnummer	NSJ-V2894-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	FACS, IF, IHC-P
Spezies Reaktivität	Human, Mouse, Rat
Immunogen	Recombinant protein was used as the immunogen for the TTF-1 antibody.
Produktbeschreibung	Recognizes a protein of 40kDa, identified as Thyroid transcription factor-1, also called NKX2.1. TTF-1 is a member of the NKx2 family of homeodomain transcription factors. It is expressed in epithelial cells of the thyroid gland and the lung. Nuclei ...
Klonalität	Monoclonal
Klon-Bezeichnung	[NX2.1/690]
UniProt	P43699
Reinheit	Protein G affinity chromatography

Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Flow cytometry: 0.5-1ug/10 ⁶ cells, Immunofluorescence: 0.5-1ug/ml, Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT
Anwendungsbeschreibung	Optimal dilution of the TTF-1 antibody should be determined by the researcher.1. Staining of formalin/paraffin tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.