

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **TTF-1 Antibody Cocktail, Clone: [8G7G3/1 + NX2.1/690], Mouse, Monoclonal NSJ-V2895IHC-7ML**

Artikelname	TTF-1 Antibody Cocktail, Clone: [8G7G3/1 + NX2.1/690], Mouse, Monoclonal
Artikelnummer	NSJ-V2895IHC-7ML
Hersteller Artikelnummer	V2895IHC-7ML
Alternativnummer	NSJ-V2895IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human, Mouse, Rat
Immunogen	Recombinant protein was used as the immunogen for the TTF-1 antibody cocktail.
Produktbeschreibung	Recognizes a protein of 40kDa, identified as Thyroid transcription factor-1, also called NKX2.1. TTF-1 is a member of the NKx2 family of homeodomain transcription factors. It is expressed in epithelial cells of the thyroid gland and the lung. Nuclei ...
Klonalität	Monoclonal
Klon-Bezeichnung	[8G7G3/1 + NX2.1/690]
UniProt	<a href="#">P43699</a>

Reinheit	Protein G affinity chromatography
Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	Optimal dilution of the TTF-1 antibody cocktail should be determined by the researcher.1. Staining of formalin/paraffin tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.