

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### TNF-alpha Antibody, Clone: [TNFA/1172], Mouse, Monoclonal NSJ-V2898IHC-7ML

Artikelname	TNF-alpha Antibody, Clone: [TNFA/1172], Mouse, Monoclonal
Artikelnummer	NSJ-V2898IHC-7ML
Hersteller Artikelnummer	V2898IHC-7ML
Alternativnummer	NSJ-V2898IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human, Rat
Immunogen	Recombinant full-length human protein was used as the immunogen for the TNF-alpha antibody.
Produktbeschreibung	This mAb recognizes human 17-26kDa protein, which is identified as cytokine TNF-alpha (Tumor Necrosis Factor-alpha). TNF-alpha can be expressed as a 17kDa free molecule, or as a 26kDa membrane protein. TNF-alpha is a protein secreted by lipopolysacch...
Klonalität	Monoclonal
Klon-Bezeichnung	[TNFA/1172]
UniProt	<a href="#">P01375</a>
Reinheit	PEG precipitation

Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	Optimal dilution of the TNF-alpha antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.