

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### TNF-alpha Antibody Cocktail, Clone: [TNF706 + P/T2], Mouse, Monoclonal NSJ-V2904-100UG

Artikelname	TNF-alpha Antibody Cocktail, Clone: [TNF706 + P/T2], Mouse, Monoclonal
Artikelnummer	NSJ-V2904-100UG
Hersteller Artikelnummer	V2904-100UG
Alternativnummer	NSJ-V2904-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	An N-terminal recombinant protein fragment (TNF706) and amino acids 115-130 (NGVELRDNQLVVPSEG) were used as the immunogen for the TNF-alpha antibody cocktail.
Produktbeschreibung	TNF alpha is a protein secreted by lipopolysaccharide-stimulated macrophages, and causes tumor necrosis when injected into tumor bearing mice. TNF alpha exists as a multimer of two, three, or five non-covalently linked units, but shows a single 17kDa...
Klonalität	Monoclonal
Klon-Bezeichnung	[TNF706 + P/T2]
UniProt	<a href="#">P01375</a>

Reinheit	PEG precipitation
Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistochemistry (FFPE): 2-4ug/ml for 30 min at RT
Anwendungsbeschreibung	Optimal dilution of the TNF-alpha antibody cocktail should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris buffer with 1mM EDTA, pH 9, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.