

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### TNF-alpha Antibody Cocktail, Clone: [TNF706 + P/T2], Mouse, Monoclonal NSJ-V2904IHC-7ML

Artikelname	TNF-alpha Antibody Cocktail, Clone: [TNF706 + P/T2], Mouse, Monoclonal
Artikelnummer	NSJ-V2904IHC-7ML
Hersteller Artikelnummer	V2904IHC-7ML
Alternativnummer	NSJ-V2904IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	An N-terminal recombinant protein fragment (TNF706) and amino acids 115-130 (NGVELRDNQLVVPSEG) were used as the immunogen for the TNF-alpha antibody cocktail.
Produktbeschreibung	TNF alpha is a protein secreted by lipopolysaccharide-stimulated macrophages, and causes tumor necrosis when injected into tumor bearing mice. TNF alpha exists as a multimer of two, three, or five non-covalently linked units, but shows a single 17kDa...
Klonalität	Monoclonal
Klon-Bezeichnung	[TNF706 + P/T2]
UniProt	<a href="#">P01375</a>

Reinheit	PEG precipitation
Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	Optimal dilution of the TNF-alpha antibody cocktail should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris buffer with 1mM EDTA, pH 9, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.