

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### TNFa Antibody, Clone: [MAb1], Mouse, Monoclonal NSJ-V2905SAF-100UG

|                          |   |
|--------------------------|---|
| Artikelname              | TNFa Antibody, Clone: [MAb1], Mouse, Monoclonal   |
| Artikelnummer            | NSJ-V2905SAF-100UG  |
| Hersteller Artikelnummer | V2905SAF-100UG  |
| Alternativnummer         | NSJ-V2905SAF-100UG  |
| Hersteller               | NSJ Bioreagents   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | FACS, IF, NeA   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant human protein was used as the immunogen for the TNFa antibody.  |
| Produktbeschreibung      | This mAb recognizes human 17-26kDa protein, which is identified as cytokine TNF-alpha (Tumor Necrosis Factor-alpha). TNF-alpha can be expressed as a 17kDa free molecule, or as a 26kDa membrane protein. TNF-alpha is a protein secreted by lipopolysacch... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [MAb1]  |
| UniProt                  | <a href="#">P01375</a>  |
| Reinheit                 | PEG precipitation   |

|                        |   |
|------------------------|---|
| Formulierung           | 1 mg/ml in 1X PBS, BSA free, sodium azide free  |
| Antibody Type          | Primary Antibody  |
| Application Verdünnung | Neutralization (order BSA/sodium azide-free format),Flow cytometry: 0.5-1ug/10 <sup>6</sup> cells,Immunofluorescence: 1-2ug/ml  |
| Anwendungsbeschreibung | Optimal dilution of the TNFa antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min. |