

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

p53 Antibody, Clone: [TRP/817], Mouse, Monoclonal NSJ-V2907SAF-100UG

| | |
|--------------------------|---|
| Artikelname | p53 Antibody, Clone: [TRP/817], Mouse, Monoclonal |
| Artikelnummer | NSJ-V2907SAF-100UG |
| Hersteller Artikelnummer | V2907SAF-100UG |
| Alternativnummer | NSJ-V2907SAF-100UG |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC-P, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human protein was used as the immunogen for the p53 antibody. |
| Produktbeschreibung | Recognizes a 53kDa protein, which is identified as p53 suppressor gene product. It reacts with the mutant as well as the wild form of p53 protein. p53 is a tumor suppressor gene expressed in a wide variety of tissue types and is involved in regulatin... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [TRP/817] |
| UniProt | P04637 |
| Reinheit | Protein G affinity chromatography |

| | |
|------------------------|--|
| Formulierung | 1 mg/ml in 1X PBS, BSA free, sodium azide free |
| Antibody Type | Primary Antibody |
| Application Verdünnung | Western blot: 0.5-1ug/ml, Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT |
| Anwendungsbeschreibung | Optimal dilution of the p53 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |