

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### GRP94 Antibody / HSP90B1, Clone: [9G10.F8.2], Rat, Monoclonal NSJ-V2908IHC-7ML

Artikelname	GRP94 Antibody / HSP90B1, Clone: [9G10.F8.2], Rat, Monoclonal
Artikelnummer	NSJ-V2908IHC-7ML
Hersteller Artikelnummer	V2908IHC-7ML
Alternativnummer	NSJ-V2908IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Rat
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human, Mouse, Rat
Immunogen	Purified glucose regulated protein 94 from chicken oviducts was used as the immunogen for the GRP94 antibody.
Produktbeschreibung	Recognizes a protein of 94kDa, which is identified as the glucose-regulated protein 94 (grp94) and also tumor rejection antigen (gp96). Grp94 shows a high degree of sequence homology with the heat shock protein 90 (hsp90). This mAb is highly specific...
Klonalität	Monoclonal
Klon-Bezeichnung	[9G10.F8.2]
UniProt	<a href="#">P14625</a>
Reinheit	Protein G affinity chromatography

Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	Optimal dilution of the GRP94 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.