

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### gp96 Antibody / GRP94 / HSP90B1, Clone: [SPM249], Rat, Monoclonal NSJ-V2909SAF-100UG

Artikelname	gp96 Antibody / GRP94 / HSP90B1, Clone: [SPM249], Rat, Monoclonal
Artikelnummer	NSJ-V2909SAF-100UG
Hersteller Artikelnummer	V2909SAF-100UG
Alternativnummer	NSJ-V2909SAF-100UG
Hersteller	NSJ Bioreagents
Wirt	Rat
Kategorie	Antikörper
Applikation	FACS, IF, IHC-P
Spezies Reaktivität	Human
Immunogen	Purified glucose regulated protein 94 from chicken oviducts was used as the immunogen for the gp96 antibody.
Produktbeschreibung	Recognizes a protein of 94kDa, which is identified as the glucose-regulated protein 94 (grp94) and also tumor rejection antigen (gp96). Gp96 shows a high degree of sequence homology with the heat shock protein 90 (hsp90). This mAb is highly specific ...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM249]
UniProt	<a href="#">P14625</a>
Reinheit	Protein G affinity chromatography

Formulierung	1 mg/ml in 1X PBS, BSA free, sodium azide free
Antibody Type	Primary Antibody
Application Verdünnung	Flow cytometry: 1-2ug/10 <sup>6</sup> cells, Immunofluorescence: 1-2ug/ml, Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	Optimal dilution of the gp96 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.