

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Tyrosinase Antibody / TYR, Clone: [OCA1/812], Mouse, Monoclonal NSJ-V2914-20UG

Artikelname	Tyrosinase Antibody / TYR, Clone: [OCA1/812], Mouse, Monoclonal
Artikelnummer	NSJ-V2914-20UG
Hersteller Artikelnummer	V2914-20UG
Alternativnummer	NSJ-V2914-20UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Recombinant human protein was used as the immunogen for the Tyrosinase antibody.
Produktbeschreibung	This mAb shows no cross-reaction with MAGE-1 and tyrosinase-related protein 1, TRP-1/gp75. Tyrosinase is a copper-containing metalloprotein that catalyzes several steps in the melanin pigment biosynthetic pathway, the hydroxylation of tyrosine t...
Klonalität	Monoclonal
Klon-Bezeichnung	[OCA1/812]
UniProt	P14679
Reinheit	Protein G affinity chromatography

Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	<p>Optimal dilution of the Tyrosinase antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9, or 10mM Citrate buffer, pH6.0, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.</p>