

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Tyrosinase Antibody / TYR, Clone: [OCA1/812], Mouse, Monoclonal NSJ-V2914IHC-7ML

Artikelname	Tyrosinase Antibody / TYR, Clone: [OCA1/812], Mouse, Monoclonal
Artikelnummer	NSJ-V2914IHC-7ML
Hersteller Artikelnummer	V2914IHC-7ML
Alternativnummer	NSJ-V2914IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Recombinant human protein was used as the immunogen for the Tyrosinase antibody.
Produktbeschreibung	This mAb shows no cross-reaction with MAGE-1 and tyrosinase-related protein 1, TRP-1/gp75. Tyrosinase is a copper-containing metalloglycoprotein that catalyzes several steps in the melanin pigment biosynthetic pathway, the hydroxylation of tyrosine t...
Klonalität	Monoclonal
Klon-Bezeichnung	[OCA1/812]
UniProt	P14679
Reinheit	Protein G affinity chromatography

Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	Optimal dilution of the Tyrosinase antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9, or 10mM Citrate buffer, pH6.0, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.