

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **TYR Antibody Cocktail (Tyrosinase), Clone: [T311 + OCA1/812], Mouse, Monoclonal NSJ-V2915IHC-7ML**

Artikelname	TYR Antibody Cocktail (Tyrosinase), Clone: [T311 + OCA1/812], Mouse, Monoclonal
Artikelnummer	NSJ-V2915IHC-7ML
Hersteller Artikelnummer	V2915IHC-7ML
Alternativnummer	NSJ-V2915IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Recombinant human protein was used as the immunogen for the TYR antibody cocktail.
Produktbeschreibung	Recognizes a cluster of proteins between 70-80kDa, identified as Tyrosinase/TYR. Occasionally a minor band at 55kDa is also detected. This mAb shows no cross-reaction with MAGE-1 and tyrosinase-related protein 1, TRP-1/gp75. Tyrosinase is a copper-co...
Klonalität	Monoclonal
Klon-Bezeichnung	[T311 + OCA1/812]
UniProt	<a href="#">P14679</a>

Reinheit	Protein G affinity chromatography
Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	Optimal dilution of the TYR antibody cocktail should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.