

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **SUMO-1 Antibody, Clone: [SUMO1/1188], Mouse, Monoclonal NSJ-V2919SAF-100UG**

Artikelname	SUMO-1 Antibody, Clone: [SUMO1/1188], Mouse, Monoclonal
Artikelnummer	NSJ-V2919SAF-100UG
Hersteller Artikelnummer	V2919SAF-100UG
Alternativnummer	NSJ-V2919SAF-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	FACS, IF, IHC-P, WB
Spezies Reaktivität	Human, Rat
Immunogen	Recombinant human protein was used as the immunogen for the SUMO-1 antibody.
Produktbeschreibung	This mAb is specific to SUMO-1 and shows no cross-reaction with either SUMO-2 or SUMO-3. The small ubiquitin-related modifier (SUMO) proteins, which include SUMO-1, SUMO-2 and SUMO-3, belong to the ubiquitin-like protein family. Like ubiquitin, the S...
Klonalität	Monoclonal
Klon-Bezeichnung	[SUMO1/1188]
UniProt	<a href="#">P63165</a>
Reinheit	Protein G affinity chromatography

Formulierung	1 mg/ml in 1X PBS, BSA free, sodium azide free
Antibody Type	Primary Antibody
Application Verdünnung	Flow cytometry: 0.5-1ug/10 <sup>6</sup> cells, Immunofluorescence: 0.5-1ug/ml, Western blot: 0.5-1ug/ml, Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT
Anwendungsbeschreibung	Optimal dilution of the SUMO-1 antibody should be determined by the researcher. 1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min. 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.