

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

VCAM-1 Antibody, Clone: [1.4C3], Mouse, Monoclonal NSJ-V2922IHC-7ML

| | |
|--------------------------|---|
| Artikelname | VCAM-1 Antibody, Clone: [1.4C3], Mouse, Monoclonal |
| Artikelnummer | NSJ-V2922IHC-7ML |
| Hersteller Artikelnummer | V2922IHC-7ML |
| Alternativnummer | NSJ-V2922IHC-7ML |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC-P |
| Spezies Reaktivität | Human |
| Immunogen | Stimulated human umbilical vein endothelial cells (HUVEC) were used as the immunogen for the VCAM-1 antibody. |
| Produktbeschreibung | Recognizes a protein of 110kDa, identified as CD106 (also known as vascular cell adhesion molecule-1 (VCAM-1), L1CAM and INCAM-100). It is a member of the Ig superfamily of adhesion molecules and is expressed at high levels on cytokine stimulated vas... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [1.4C3] |
| UniProt | P19320 |
| Reinheit | Protein G affinity chromatography |

| | |
|------------------------|---|
| Formulierung | Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only* |
| Antibody Type | Primary Antibody |
| Application Verdünnung | The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |
| Anwendungsbeschreibung | Optimal dilution of the VCAM-1 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |