

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Vimentin Antibody, Clone: [LN-6], Mouse, Monoclonal NSJ-V2927-20UG

Artikelname	Vimentin Antibody, Clone: [LN-6], Mouse, Monoclonal
Artikelnummer	NSJ-V2927-20UG
Hersteller Artikelnummer	V2927-20UG
Alternativnummer	NSJ-V2927-20UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human, Mouse, Rat
Immunogen	A human thymic nuclear extract was used as the immunogen for the Vimentin antibody.
Produktbeschreibung	This mAb reacts with a 58kDa protein identified as vimentin. It reacts with a non-hematopoietic epitope of vimentin and shows no cross-reaction with other closely related intermediate filament proteins (IFPs) such as desmin, keratin, neurofilament, a...
Klonalität	Monoclonal
Klon-Bezeichnung	[LN-6]
UniProt	<a href="#">P08670</a>
Reinheit	PEG precipitation

Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistochemistry (FFPE): 0.1-0.2ug/ml for 30 min at RT
Anwendungsbeschreibung	Optimal dilution of the Vimentin antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.