

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Anti-Vimentin Antibody, Clone: [VM1170], Mouse, Monoclonal NSJ-V2928-20UG

Artikelname	Anti-Vimentin Antibody, Clone: [VM1170], Mouse, Monoclonal
Artikelnummer	NSJ-V2928-20UG
Hersteller Artikelnummer	V2928-20UG
Alternativnummer	NSJ-V2928-20UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P, WB
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human protein was used as the immunogen for the anti-Vimentin antibody.
Produktbeschreibung	This mAb reacts with a 58kDa protein identified as vimentin. It shows no cross-reaction with other closely related intermediate filament proteins (IFPs) such as desmin, keratin, neurofilament, and glial fibrillary acid protein. Anti-vimentin alone is...
Klonalität	Monoclonal
Klon-Bezeichnung	[VM1170]
UniProt	<a href="#">P08670</a>
Reinheit	Protein G affinity chromatography

Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Western blot: 1-2ug/ml, Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	Optimal dilution of the anti-Vimentin antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.