

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Anti-vWF Antibody, Clone: [IIIE2.34], Mouse, Monoclonal NSJ-V2929-100UG

| | |
|--------------------------|---|
| Artikelname | Anti-vWF Antibody, Clone: [IIIE2.34], Mouse, Monoclonal |
| Artikelnummer | NSJ-V2929-100UG |
| Hersteller Artikelnummer | V2929-100UG |
| Alternativnummer | NSJ-V2929-100UG |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC-P |
| Spezies Reaktivität | Human |
| Immunogen | A recombinant human protein fragment spanning aa 845-949 was used as the immunogen for the anti-vWF antibody. |
| Produktbeschreibung | von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslation... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [IIIE2.34] |
| UniProt | P04275 |
| Reinheit | Protein G affinity chromatography |

| | |
|------------------------|---|
| Formulierung | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide |
| Antibody Type | Primary Antibody |
| Application Verdünnung | Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT |
| Anwendungsbeschreibung | Optimal dilution of the anti-vWF antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min. |