

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

WT1 Antibody, Clone: [WT1/857], Mouse, Monoclonal NSJ-V2932IHC-7ML

| | |
|--------------------------|---|
| Artikelname | WT1 Antibody, Clone: [WT1/857], Mouse, Monoclonal |
| Artikelnummer | NSJ-V2932IHC-7ML |
| Hersteller Artikelnummer | V2932IHC-7ML |
| Alternativnummer | NSJ-V2932IHC-7ML |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC-P |
| Spezies Reaktivität | Human, Rat |
| Immunogen | A recombinant human protein was used as the immunogen for the WT1 antibody. |
| Produktbeschreibung | Recognizes a 47-55kDa-tumor suppressor protein, identified as Wilms Tumor (WT1) protein. The antibody reacts with all isoforms of the full-length WT1 and also identifies WT1 lacking exon 2-encoded amino acids, frequently found in subsets of sporadic ... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [WT1/857] |
| UniProt | P19544 |
| Reinheit | Protein G affinity chromatography |

| | |
|------------------------|--|
| Formulierung | Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only* |
| Antibody Type | Primary Antibody |
| Application Verdünnung | The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |
| Anwendungsbeschreibung | Optimal dilution of the WT1 antibody should be determined by the researcher.1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. 2. View the recombinant version of the WT1/857 mAb. |