

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

WT1 Antibody Cocktail, Clone: [WT1/857 + 6F-H2], Mouse, Monoclonal NSJ-V2933-100UG

Artikelname	WT1 Antibody Cocktail, Clone: [WT1/857 + 6F-H2], Mouse, Monoclonal
Artikelnummer	NSJ-V2933-100UG
Hersteller Artikelnummer	V2933-100UG
Alternativnummer	NSJ-V2933-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Recombinant full length human protein (WT1/857) and amino acids 1-181 of human Wilms Tumor (6F-H2) were used as the immunogen for the WT1 antibody cocktail.
Produktbeschreibung	Recognizes a 47-55kDa-tumor suppressor protein, identified as Wilms Tumor (WT1) protein. The antibody reacts with all isoforms of the full-length WT1 and also identifies WT1 lacking exon 2-encoded amino acids, frequently found in subsets of sporadic ...
Klonalität	Monoclonal
Klon-Bezeichnung	[WT1/857 + 6F-H2]
UniProt	P19544

Reinheit	Protein G affinity chromatography
Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	Optimal dilution of the WT1 antibody cocktail should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.