

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

WT1 Antibody Cocktail, Clone: [WT1/857 + 6F-H2], Mouse, Monoclonal NSJ-V2933IHC-7ML

Artikelname	WT1 Antibody Cocktail, Clone: [WT1/857 + 6F-H2], Mouse, Monoclonal
Artikelnummer	NSJ-V2933IHC-7ML
Hersteller Artikelnummer	V2933IHC-7ML
Alternativnummer	NSJ-V2933IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Recombinant full length human protein (WT1/857) and amino acids 1-181 of human Wilms Tumor (6F-H2) were used as the immunogen for the WT1 antibody cocktail.
Produktbeschreibung	Recognizes a 47-55kDa-tumor suppressor protein, identified as Wilms Tumor (WT1) protein. The antibody reacts with all isoforms of the full-length WT1 and also identifies WT1 lacking exon 2-encoded amino acids, frequently found in subsets of sporadic ...
Klonalität	Monoclonal
Klon-Bezeichnung	[WT1/857 + 6F-H2]
UniProt	P19544

Reinheit	Protein G affinity chromatography
Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	Optimal dilution of the WT1 antibody cocktail should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.