

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

ZAP-70 Antibody, Clone: [2F3.2], Mouse, Monoclonal NSJ-V2934SAF-100UG

| | |
|--------------------------|---|
| Artikelname | ZAP-70 Antibody, Clone: [2F3.2], Mouse, Monoclonal |
| Artikelnummer | NSJ-V2934SAF-100UG |
| Hersteller Artikelnummer | V2934SAF-100UG |
| Alternativnummer | NSJ-V2934SAF-100UG |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FACS, IF, IHC-P, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human protein (including amino acids 1-254 and the SH2 domains) was used as the immunogen for the ZAP-70 antibody. |
| Produktbeschreibung | ZAP70 is a 70kDa protein tyrosine kinase found in T-cells and natural killer cells. Control of this protein translation is via the IgVH gene. ZAP70 protein is expressed in leukemic cells of approximately 25% of chronic lymphocytic leukemia (CLL) case... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [2F3.2] |
| UniProt | P43403 |
| Reinheit | Protein G affinity chromatography |

| | |
|------------------------|---|
| Formulierung | 1 mg/ml in 1X PBS, BSA free, sodium azide free |
| Antibody Type | Primary Antibody |
| Application Verdünnung | Western blot: 1-2ug/ml,Flow cytometry: 1-2ug/10 ⁶ cells,Immunofluorescence: 1-2ug/ml,Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT |
| Anwendungsbeschreibung | Optimal dilution of the ZAP-70 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |