

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### ZAP70 Antibody Cocktail, Clone: [ZAP70/528 + 2F3.2], Mouse, Monoclonal NSJ-V2937SAF-100UG

Artikelname	ZAP70 Antibody Cocktail, Clone: [ZAP70/528 + 2F3.2], Mouse, Monoclonal
Artikelnummer	NSJ-V2937SAF-100UG
Hersteller Artikelnummer	V2937SAF-100UG
Alternativnummer	NSJ-V2937SAF-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human protein was used as the immunogen for the ZAP70 antibody cocktail.
Produktbeschreibung	ZAP70 is a 70kDa protein tyrosine kinase found in T-cells and natural killer cells. Control of this protein translation is via the IgVH gene. ZAP70 protein is expressed in leukemic cells of approximately 25% of chronic lymphocytic leukemia (CLL) case...
Klonalität	Monoclonal
Klon-Bezeichnung	[ZAP70/528 + 2F3.2]
UniProt	<a href="#">P43403</a>
Reinheit	Protein G affinity chromatography

Formulierung	1 mg/ml in 1X PBS, BSA free, sodium azide free
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	<p>Optimal dilution of the ZAP70 antibody cocktail should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.</p>