

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Caldesmon Antibody / CALD1 / CAD (HMW), Clone: [SPM168], Mouse, Monoclonal NSJ-V2941IHC-7ML

| | |
|--------------------------|---|
| Artikelname | Caldesmon Antibody / CALD1 / CAD (HMW), Clone: [SPM168], Mouse, Monoclonal |
| Artikelnummer | NSJ-V2941IHC-7ML |
| Hersteller Artikelnummer | V2941IHC-7ML |
| Alternativnummer | NSJ-V2941IHC-7ML |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC-P |
| Spezies Reaktivität | Human |
| Immunogen | Crude human uterus extract was used as the immunogen for the Caldesmon antibody. |
| Produktbeschreibung | Recognizes a protein of 150kDa, which is identified as the high molecular weight variant of Caldesmon. Two closely related variants of human caldesmon have been identified which are different in their electrophoretic mobility and cellular distributio... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [SPM168] |
| UniProt | Q05682 |

| | |
|------------------------|--|
| Reinheit | Protein G affinity chromatography |
| Formulierung | Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only* |
| Antibody Type | Primary Antibody |
| Application Verdünnung | The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |
| Anwendungsbeschreibung | Optimal dilution of the Caldesmon antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |