

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD30 Antibody, Clone: [Ber-H2], Mouse, Monoclonal NSJ-V2990S-0.5ML

| | |
|--------------------------|---|
| Artikelname | CD30 Antibody, Clone: [Ber-H2], Mouse, Monoclonal |
| Artikelnummer | NSJ-V2990S-0.5ML |
| Hersteller Artikelnummer | V2990S-0.5ML |
| Alternativnummer | NSJ-V2990S-0.5ML |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC-P |
| Spezies Reaktivität | Human |
| Immunogen | A Co cell line established from a patient with Hodgkin s disease of T-cell lineage was used as the immunogen for the CD30 antibody. |
| Produktbeschreibung | Recognizes a single chain glycoprotein of 105/120kDa, identified as CD30/Ki-1. Its epitope is located between aa112-412. CD30 is synthesized as a 90kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 105... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [Ber-H2] |
| UniProt | P28908 |
| Reinheit | Unpurified high titer supernatant |

| | |
|------------------------|---|
| Formulierung | Bioreactor concentrate with 0.05% sodium azide |
| Antibody Type | Primary Antibody |
| Application Verdünnung | Immunohistochemistry (FFPE): 1:50-1:100 for 30 min at RT (1) |
| Anwendungsbeschreibung | Optimal dilution of the CD30 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 min. |