

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD30 Antibody, Clone: [Ki-1/779], Mouse, Monoclonal NSJ-V2992SAF-100UG

Artikelname	CD30 Antibody, Clone: [Ki-1/779], Mouse, Monoclonal
Artikelnummer	NSJ-V2992SAF-100UG
Hersteller Artikelnummer	V2992SAF-100UG
Alternativnummer	NSJ-V2992SAF-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	FACS, IF, IHC-P
Spezies Reaktivität	Human
Immunogen	Recombinant human protein was used as the immunogen for the CD30 antibody.
Produktbeschreibung	CD30 is the receptor for TNFSF8/CD30L. May play a role in the regulation of cellular growth and transformation of activated lymphoblasts. Regulates gene expression through activation of NF-kappa-B. [UniProt]...
Klonalität	Monoclonal
Klon-Bezeichnung	[Ki-1/779]
UniProt	P28908
Reinheit	Protein G affinity chromatography

Formulierung	1 mg/ml in 1X PBS, BSA free, sodium azide free
Antibody Type	Primary Antibody
Application Verdünnung	Flow cytometry: 0.5-1ug/10 ⁶ cells, Immunofluorescence: 1-2ug/ml, Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT
Anwendungsbeschreibung	Optimal dilution of the CD30 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.