

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Napsin A Antibody, Clone: [NAPSA/1239], Mouse, Monoclonal NSJ-V2996-100UG

Artikelname	Napsin A Antibody, Clone: [NAPSA/1239], Mouse, Monoclonal
Artikelnummer	NSJ-V2996-100UG
Hersteller Artikelnummer	V2996-100UG
Alternativnummer	NSJ-V2996-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	FACS, IF, IHC-P, WB
Spezies Reaktivität	Human
Immunogen	A recombinant fragment from the human protein (amino acids 189-299) was used as the immunogen for the Napsin A antibody.
Produktbeschreibung	Napsin-A is a protein that in humans is encoded by the NAPSA gene. The activation peptides of aspartic proteinases plays role as inhibitors of the active site. These peptide segments, or pro-parts, are deemed important for correct folding, targeting,...
Klonalität	Monoclonal
Klon-Bezeichnung	[NAPSA/1239]
UniProt	O96009
Reinheit	Protein G affinity chromatography

Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Flow cytometry: 0.5-1ug/10 ⁶ cells, Immunofluorescence: 1-2ug/ml, Western blot: 1-2ug/ml for 60 min at RT, Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	Optimal dilution of the Napsin A antibody should be determined by the researcher. 1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min. 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.