

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Napsin A Antibody Cocktail, Clone: [NAPSA/1238 + NAPSA/1239], Mouse, Monoclonal NSJ-V2997IHC-7ML

Artikelname	Napsin A Antibody Cocktail, Clone: [NAPSA/1238 + NAPSA/1239], Mouse, Monoclonal
Artikelnummer	NSJ-V2997IHC-7ML
Hersteller Artikelnummer	V2997IHC-7ML
Alternativnummer	NSJ-V2997IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Amino acids 189-299 from the human protein were used as the immunogen for the mAbs in the Napsin A antibody cocktail.
Produktbeschreibung	Napsin is a pepsin-like aspartic proteinase connected with maturation of surfactant protein B. There are two closely related napsins, napsin A and napsin B. Napsin A is expressed as a single chain protein. Immunohistochemical studies revealed high ex...
Klonalität	Monoclonal
Klon-Bezeichnung	[NAPSA/1238 + NAPSA/1239]
UniProt	O96009

Reinheit	Protein G affinity chromatography
Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	Optimal dilution of the Napsin A antibody cocktail should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.