

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### CD44 Antibody, Clone: [DF1485], Mouse, Monoclonal NSJ-V3012-100UG

Artikelname	CD44 Antibody, Clone: [DF1485], Mouse, Monoclonal
Artikelnummer	NSJ-V3012-100UG
Hersteller Artikelnummer	V3012-100UG
Alternativnummer	NSJ-V3012-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IF, IHC-P, WB
Spezies Reaktivität	Human, Mouse
Immunogen	Purified CD44 antigen (PGp-1) from lymphocyte membrane was used as the immunogen for the CD44 antibody.
Produktbeschreibung	Recognizes a cell surface glycoprotein of 80-95kDa (CD44/HCAM) on lymphocytes, monocytes, and granulocytes (Leucocyte Typing Workshop V). Its epitope is resistant to digestion by trypsin and chymotrypsin. The CD44 family of glycoproteins exists in a ...
Klonalität	Monoclonal
Klon-Bezeichnung	[DF1485]
UniProt	<a href="#">P16070</a>
Reinheit	Protein G affinity chromatography

Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Western blot: 1-2ug/ml, Immunofluorescence: 1-2ug/ml, Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	Optimal dilution of the CD44 antibody should be determined by the researcher.1. Staining of formalin/paraffin tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.