

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### CD44 Antibody, Clone: [DF1485], Mouse, Monoclonal NSJ-V3012IHC-7ML

Artikelname	CD44 Antibody, Clone: [DF1485], Mouse, Monoclonal
Artikelnummer	NSJ-V3012IHC-7ML
Hersteller Artikelnummer	V3012IHC-7ML
Alternativnummer	NSJ-V3012IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human, Mouse
Immunogen	Purified CD44 antigen (PGp-1) from lymphocyte membrane was used as the immunogen for the CD44 antibody.
Produktbeschreibung	Recognizes a cell surface glycoprotein of 80-95kDa (CD44/HCAM) on lymphocytes, monocytes, and granulocytes (Leucocyte Typing Workshop V). Its epitope is resistant to digestion by trypsin and chymotrypsin. The CD44 family of glycoproteins exists in a ...
Klonalität	Monoclonal
Klon-Bezeichnung	[DF1485]
UniProt	<a href="#">P16070</a>
Reinheit	Protein G affinity chromatography

Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	Optimal dilution of the CD44 antibody should be determined by the researcher.1. Staining of formalin/paraffin tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.