

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD59 Antibody, Clone: [MACIF/1193], Mouse, Monoclonal NSJ-V3024IHC-7ML

| | |
|--------------------------|---|
| Artikelname | CD59 Antibody, Clone: [MACIF/1193], Mouse, Monoclonal |
| Artikelnummer | NSJ-V3024IHC-7ML |
| Hersteller Artikelnummer | V3024IHC-7ML |
| Alternativnummer | NSJ-V3024IHC-7ML |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC-P |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length human protein was used as the immunogen for the CD59 antibody. |
| Produktbeschreibung | Reacts with human CD59, a 20kDa glycosyl phosphatidyl-inositol (GPI)-anchored cell surface protein. CD59 regulates complement-mediated cell lysis, and it is involved in lymphocyte signal transduction. This protein is a potent inhibitor of the complem... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [MACIF/1193] |
| UniProt | P13987 |
| Reinheit | PEG precipitation |

| | |
|------------------------|---|
| Formulierung | Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only* |
| Antibody Type | Primary Antibody |
| Application Verdünnung | The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |
| Anwendungsbeschreibung | Optimal dilution of the CD59 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |