

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

LAMP-3 Antibody / CD63, Clone: [LAMP3/968], Mouse, Monoclonal NSJ-V3029-20UG

Artikelname	LAMP-3 Antibody / CD63, Clone: [LAMP3/968], Mouse, Monoclonal
Artikelnummer	NSJ-V3029-20UG
Hersteller Artikelnummer	V3029-20UG
Alternativnummer	NSJ-V3029-20UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	FACS, IHC-P
Spezies Reaktivität	Human
Immunogen	Recombinant human full-length protein was used as the immunogen for the CD63 / LAMP-3 antibody.
Produktbeschreibung	This mAb recognizes protein of 26kDa-60kDa, which is identified as CD63/LAMP-3. Its epitope is different from that of mAb LAMP3/529. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cel...
Klonalität	Monoclonal
Klon-Bezeichnung	[LAMP3/968]
UniProt	P08962
Reinheit	Protein G affinity chromatography

Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Flow cytometry: 1-2ug/10 ⁶ cells, Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	Optimal dilution of the LAMP-3 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.