

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD63 Antibody Cocktail, Clone: [NKI/C3 + LAMP3/968], Mouse, Monoclonal NSJ-V3030-100UG

| | |
|--------------------------|---|
| Artikelname | CD63 Antibody Cocktail, Clone: [NKI/C3 + LAMP3/968], Mouse, Monoclonal |
| Artikelnummer | NSJ-V3030-100UG |
| Hersteller Artikelnummer | V3030-100UG |
| Alternativnummer | NSJ-V3030-100UG |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FACS, IF, IHC-P |
| Spezies Reaktivität | Human |
| Immunogen | The smooth plasma membrane fraction of MeWo cells (NKI/C3) and recombinant human full-length protein (LAMP3/968) were used as the immunogen for the CD63 antibody cocktail. |
| Produktbeschreibung | This mAb recognizes protein of 26kDa-60kDa, which is identified as CD63 / LAMP-3. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multi-molecular complexes ... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [NKI/C3 + LAMP3/968] |
| UniProt | P08962 |

| | |
|------------------------|--|
| Reinheit | Protein G affinity chromatography |
| Formulierung | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide |
| Antibody Type | Primary Antibody |
| Application Verdünnung | Flow cytometry: 0.5-1ug/10 ⁶ cells, Immunofluorescence: 0.5-1ug/ml, Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT |
| Anwendungsbeschreibung | Optimal dilution of the CD63 antibody cocktail should be determined by the researcher. 1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min. 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |