

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### CD63 Antibody Cocktail, Clone: [NKI/C3 + LAMP3/968], Mouse, Monoclonal NSJ-V3030-20UG

Artikelname	CD63 Antibody Cocktail, Clone: [NKI/C3 + LAMP3/968], Mouse, Monoclonal
Artikelnummer	NSJ-V3030-20UG
Hersteller Artikelnummer	V3030-20UG
Alternativnummer	NSJ-V3030-20UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	FACS, IF, IHC-P
Spezies Reaktivität	Human
Immunogen	The smooth plasma membrane fraction of MeWo cells (NKI/C3) and recombinant human full-length protein (LAMP3/968) were used as the immunogen for the CD63 antibody cocktail.
Produktbeschreibung	This mAb recognizes protein of 26kDa-60kDa, which is identified as CD63 / LAMP-3. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multi-molecular complexes ...
Klonalität	Monoclonal
Klon-Bezeichnung	[NKI/C3 + LAMP3/968]
UniProt	<a href="#">P08962</a>

Reinheit	Protein G affinity chromatography
Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Flow cytometry: 0.5-1ug/10 <sup>6</sup> cells, Immunofluorescence: 0.5-1ug/ml, Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	Optimal dilution of the CD63 antibody cocktail should be determined by the researcher. 1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min. 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.