

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD68 Antibody, Clone: [LAMP4/824], Mouse, Monoclonal NSJ-V3031-20UG

Artikelname	CD68 Antibody, Clone: [LAMP4/824], Mouse, Monoclonal
Artikelnummer	NSJ-V3031-20UG
Hersteller Artikelnummer	V3031-20UG
Alternativnummer	NSJ-V3031-20UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	FACS, IHC-P, WB
Spezies Reaktivität	Human
Immunogen	Recombinant human protein was used as the immunogen for the CD68 antibody.
Produktbeschreibung	CD68 could play a role in phagocytic activities of tissue macrophages, both in intracellular lysosomal metabolism and extracellular cell-cell and cell-pathogen interactions. Binds to tissue- and organ-specific lectins or selectins, allowing homing of...
Klonalität	Monoclonal
Klon-Bezeichnung	[LAMP4/824]
UniProt	P34810
Reinheit	Protein G affinity chromatography

Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Western blot: 1-2ug/ml,Flow cytometry: 1-2ug/10 ⁶ cells,Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT
Anwendungsbeschreibung	Optimal dilution of the CD68 antibody should be determined by the researcher.1. Staining of formalin/paraffin tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.