

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD70 Antibody, Clone: [BU69], Mouse, Monoclonal NSJ-V3033-20UG

| | |
|--------------------------|---|
| Artikelname | CD70 Antibody, Clone: [BU69], Mouse, Monoclonal |
| Artikelnummer | NSJ-V3033-20UG |
| Hersteller Artikelnummer | V3033-20UG |
| Alternativnummer | NSJ-V3033-20UG |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FACS, IF, IHC |
| Spezies Reaktivität | Human |
| Immunogen | The human WM-1 (Waldenstrom s macroglobulinemia) cell line was used as the immunogen for the CD70 antibody. |
| Produktbeschreibung | CD70 belongs to the tumor necrosis factor (TNF) ligand family. It is a cytokine that binds to CD27. Plays a role in T-cell activation. Induces the proliferation of costimulated T-cells and enhances the generation of cytolytic T-cells.... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [BU69] |
| UniProt | P32970 |
| Reinheit | Protein G affinity chromatography |

| | |
|------------------------|---|
| Formulierung | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide |
| Antibody Type | Primary Antibody |
| Application Verdünnung | Flow cytometry: 1-2ug/million cells, Immunofluorescence: 1-3ug/ml, Immunohistochemistry: 1-2ug/ml for 30 min at RT |
| Anwendungsbeschreibung | Optimal dilution of the CD70 antibody should be determined by the researcher.1. This mAb blocks the interaction between CD27 and CD70, and has been shown to inhibit T cell proliferation induced by dendritic cells. |