

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cdc20 Antibody, Clone: [CDC20/1102], Mouse, Monoclonal NSJ-V3041-20UG

| | |
|--------------------------|---|
| Artikelname | Cdc20 Antibody, Clone: [CDC20/1102], Mouse, Monoclonal |
| Artikelnummer | NSJ-V3041-20UG |
| Hersteller Artikelnummer | V3041-20UG |
| Alternativnummer | NSJ-V3041-20UG |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC-P |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human protein was used as the immunogen for the Cdc20 antibody. |
| Produktbeschreibung | Cyclins, regulatory subunits, which associate with kinases, control many of the important steps in cell cycle progression. The Cdc2 protein kinase (p34Cdc2) exhibits protein kinase activity in vitro and exists in a complex with both cyclin B and a pr... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [CDC20/1102] |
| UniProt | Q12834 |
| Reinheit | Protein G affinity chromatography |

| | |
|------------------------|--|
| Formulierung | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide |
| Antibody Type | Primary Antibody |
| Application Verdünnung | Immunohistochemistry (FFPE): 2-4ug/ml for 30 min at RT |
| Anwendungsbeschreibung | Optimal dilution of the Cdc20 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |