

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Recombinant Cytokeratin 10 Antibody / Rabbit Monoclonal, Clone: [CTKN10-2R] NSJ-V3711SAF-100UG

|                          |   |
|--------------------------|---|
| Artikelname              | Recombinant Cytokeratin 10 Antibody / Rabbit Monoclonal, Clone:<br>[CTKN10-2R]  |
| Artikelnummer            | NSJ-V3711SAF-100UG  |
| Hersteller Artikelnummer | V3711SAF-100UG  |
| Alternativnummer         | NSJ-V3711SAF-100UG  |
| Hersteller               | NSJ Bioreagents   |
| Wirt                     | Rabbit  |
| Kategorie                | Antikörper  |
| Applikation              | IHC-P   |
| Spezies Reaktivität      | Human   |
| Immunogen                | A cytoskeletal preparation extracted from human ectocervical epithelium was used as the immunogen for the recombinant Cytokeratin 10 antibody.  |
| Produktbeschreibung      | Cytokeratin-10 (CK-10) or keratin-10 (K10) is a protein that in humans is encoded by the KRT10 gene. CK10 is a member of the type I (acidic) cytokeratin family, which belongs to the superfamily of intermediate filament (IF) proteins. Keratins are het... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [CTKN10-2R]   |
| UniProt                  | <a href="#">P13645</a>  |

|                        |  |
|------------------------|--|
| Reinheit               | Protein A affinity chromatography  |
| Formulierung           | 1 mg/ml in 1X PBS, BSA free, sodium azide free   |
| Antibody Type          | Primary Antibody   |
| Application Verdünnung | Immunohistochemistry (FFPE): 0.1-0.2ug/ml for 30 min at RT, Prediluted IHC only format: incubate for 30 min at RT (1)  |
| Anwendungsbeschreibung | The stated application concentrations are suggested starting points. Titration of the recombinant Cytokeratin 10 antibody may be required due to differences in protocols and secondary/substrate sensitivity.1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |