

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Recombinant GPC3 Antibody / Glypican-3, Clone: [rGPC3/863], Mouse, Monoclonal NSJ-V3714-100UG

|                          |  |
|--------------------------|--|
| Artikelname              | Recombinant GPC3 Antibody / Glypican-3, Clone: [rGPC3/863], Mouse, Monoclonal  |
| Artikelnummer            | NSJ-V3714-100UG  |
| Hersteller Artikelnummer | V3714-100UG  |
| Alternativnummer         | NSJ-V3714-100UG  |
| Hersteller               | NSJ Bioreagents  |
| Wirt                     | Mouse  |
| Kategorie                | Antikörper   |
| Applikation              | FACS, IF, IHC-P  |
| Spezies Reaktivität      | Human  |
| Immunogen                | Recombinant full-length human Glypican 3 protein was used as the immunogen for the recombinant GPC3 antibody.  |
| Produktbeschreibung      | Glypican-3 (GPC3) is a glycosylphosphatidyl inositol-anchored membrane protein, which may also be found in a secreted form. Anti-GPC3 has been identified as a useful tumor marker for the diagnosis of hepatocellular carcinoma (HCC), hepatoblastoma, mel... |
| Klonalität               | Monoclonal   |
| Klon-Bezeichnung         | [rGPC3/863]  |
| UniProt                  | <a href="#">P51654</a>   |

|                        |  |
|------------------------|--|
| Reinheit               | Protein G affinity chromatography  |
| Formulierung           | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide   |
| Antibody Type          | Primary Antibody   |
| Application Verdünnung | Flow cytometry: 1-2ug/million cells, Immunofluorescence: 1-2ug/ml, Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT  |
| Anwendungsbeschreibung | <p>Optimal dilution of the recombinant GPC3 antibody should be determined by the researcher.1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.&lt;br /&gt;2. View the non-recombinant version of this &lt;a href=../tds/glypican-3-antibody-gpc3-863-v2538&gt;GPC3 antibody.</p> |