

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Recombinant Multi Cytokeratin Antibody / Rabbit Monoclonal, Clone: [CTKNM-1R] NSJ-V3718-100UG

| | |
|--------------------------|--|
| Artikelname | Recombinant Multi Cytokeratin Antibody / Rabbit Monoclonal, Clone: [CTKNM-1R] |
| Artikelnummer | NSJ-V3718-100UG |
| Hersteller Artikelnummer | V3718-100UG |
| Alternativnummer | NSJ-V3718-100UG |
| Hersteller | NSJ Bioreagents |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IF, IHC-P |
| Spezies Reaktivität | Human |
| Immunogen | Human epithelial cell keratin was used as the immunogen for the recombinant Multi Cytokeratin antibody. |
| Produktbeschreibung | Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pI 6.0) subfamilies. This mAb recognizes acidic (Type I or LMW) and basic (Type II or HMW) cytokeratins, including 59kDa (CK4), 58kDa (CK5), 56kDa (CK6), 52kDa (...) |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [CTKNM-1R] |
| Reinheit | Protein A affinity chromatography |

| | |
|------------------------|---|
| Formulierung | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide |
| Antibody Type | Primary Antibody |
| Application Verdünnung | Immunofluorescence: 1-2ug/ml, Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT, Prediluted IHC only format: incubate for 30 min at RT (1) |
| Anwendungsbeschreibung | Optimal dilution of the recombinant Multi Cytokeratin antibody should be determined by the researcher. 1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |