

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Recombinant Multi Cytokeratin Antibody / Rabbit Monoclonal, Clone: [CTKNM-1R] NSJ-V3718-20UG

Artikelname	Recombinant Multi Cytokeratin Antibody / Rabbit Monoclonal, Clone: [CTKNM-1R]
Artikelnummer	NSJ-V3718-20UG
Hersteller Artikelnummer	V3718-20UG
Alternativnummer	NSJ-V3718-20UG
Hersteller	NSJ Bioreagents
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IF, IHC-P
Spezies Reaktivität	Human
Immunogen	Human epithelial cell keratin was used as the immunogen for the recombinant Multi Cytokeratin antibody.
Produktbeschreibung	Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pI 6.0) subfamilies. This mAb recognizes acidic (Type I or LMW) and basic (Type II or HMW) cytokeratins, including 59kDa (CK4), 58kDa (CK5), 56kDa (CK6), 52kDa (...)
Klonalität	Monoclonal
Klon-Bezeichnung	[CTKNM-1R]
Reinheit	Protein A affinity chromatography

Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Immunofluorescence: 1-2ug/ml, Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT, Prediluted IHC only format: incubate for 30 min at RT (1)
Anwendungsbeschreibung	Optimal dilution of the recombinant Multi Cytokeratin antibody should be determined by the researcher. 1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.