

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### NKX2.2 Antibody, Clone: [RMNK2-2], Mouse, Monoclonal NSJ-V3726SAF-100UG

|                          |   |
|--------------------------|---|
| Artikelname              | NKX2.2 Antibody, Clone: [RMNK2-2], Mouse, Monoclonal  |
| Artikelnummer            | NSJ-V3726SAF-100UG  |
| Hersteller Artikelnummer | V3726SAF-100UG  |
| Alternativnummer         | NSJ-V3726SAF-100UG  |
| Hersteller               | NSJ Bioreagents   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC-P   |
| Spezies Reaktivität      | Human   |
| Immunogen                | A human partial recombinant protein corresponding to amino acids 1-119 was used as the immunogen for this NKX2.2 antibody.  |
| Produktbeschreibung      | Expression of NKX2.2 has been found in neuroendocrine tumors of the gut, making it a potential marker for the study of gastrointestinal neuroendocrine tumors. More recently, NKX2.2 protein was identified as a target of EWS-FLI-1, the fusion protein s... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [RMNK2-2]   |
| UniProt                  | <a href="#">O95096</a>  |
| Reinheit                 | Protein G affinity chromatography   |

|                        |  |
|------------------------|--|
| Formulierung           | 1 mg/ml in 1X PBS, BSA free, sodium azide free   |
| Antibody Type          | Primary Antibody   |
| Application Verdünnung | Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT, Prediluted IHC only format : incubate for 30 min at RT (1)   |
| Anwendungsbeschreibung | Titering of the NKX2.2 antibody may be required for optimal performance.1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |