

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Recombinant CD20 Antibody, Clone: [rIGEL/773], Mouse, Monoclonal NSJ-V3727SAF-100UG

| | |
|--------------------------|---|
| Artikelname | Recombinant CD20 Antibody, Clone: [rIGEL/773], Mouse, Monoclonal |
| Artikelnummer | NSJ-V3727SAF-100UG |
| Hersteller Artikelnummer | V3727SAF-100UG |
| Alternativnummer | NSJ-V3727SAF-100UG |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FACS, IF, IHC-P, WB |
| Spezies Reaktivität | Human |
| Immunogen | Human CD20 protein was used as the immunogen for this recombinant CD20 antibody. |
| Produktbeschreibung | Recognizes a protein of 30-33kDa, which is identified as CD20. It is a non-Ig differentiation antigen of B-cells and its expression is restricted to normal and neoplastic B-cells, being absent from all other leukocytes and tissues. CD20 is expressed ... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [rIGEL/773] |
| UniProt | P11836 |
| Reinheit | Protein G affinity chromatography |

| | |
|------------------------|--|
| Formulierung | 1 mg/ml in 1X PBS, BSA free, sodium azide free |
| Antibody Type | Primary Antibody |
| Application Verdünnung | Immunofluorescence: 1-3ug/ml,Flow cytometry: 1-2ug/million cells,Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT,Western blot: 2-4ug/ml |
| Anwendungsbeschreibung | Titering of the recombinant CD20 antibody may be required for optimal performance.1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |