

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Recombinant Thyroglobulin Antibody / Rabbit Monoclonal, Clone: [TG157-3R] NSJ-V3734-100UG

| | |
|--------------------------|--|
| Artikelname | Recombinant Thyroglobulin Antibody / Rabbit Monoclonal, Clone: [TG157-3R] |
| Artikelnummer | NSJ-V3734-100UG |
| Hersteller Artikelnummer | V3734-100UG |
| Alternativnummer | NSJ-V3734-100UG |
| Hersteller | NSJ Bioreagents |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC-P |
| Immunogen | Human thyroid follicular cells were used as the immunogen for the recombinant Thyroglobulin antibody. |
| Produktbeschreibung | Thyroglobulin (Tg) is a glycoprotein homodimer produced predominantly by the thyroid gland. It acts as a substrate for the synthesis of thyroxine and triiodothyronine as well as the storage of the inactive forms of thyroid hormone and iodine. Thyrog... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [TG157-3R] |
| UniProt | P01266 |
| Reinheit | Protein A affinity chromatography |

| | |
|------------------------|--|
| Formulierung | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide |
| Antibody Type | Primary Antibody |
| Application Verdünnung | Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT, Prediluted IHC only format: incubate for 30 min at RT (1) |
| Anwendungsbeschreibung | Optimal dilution of the recombinant Thyroglobulin antibody should be determined by the researcher.1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |