

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Recombinant ATRX Antibody / RAD54, Clone: [ATMRX-1R], Rabbit, Monoclonal NSJ-V3991SAF-100UG

|                          |   |
|--------------------------|---|
| Artikelname              | Recombinant ATRX Antibody / RAD54, Clone: [ATMRX-1R], Rabbit, Monoclonal  |
| Artikelnummer            | NSJ-V3991SAF-100UG  |
| Hersteller Artikelnummer | V3991SAF-100UG  |
| Alternativnummer         | NSJ-V3991SAF-100UG  |
| Hersteller               | NSJ Bioreagents   |
| Wirt                     | Rabbit  |
| Kategorie                | Antikörper  |
| Applikation              | IHC-P   |
| Immunogen                | Recombinant human ATRX protein was used as the immunogen for this recombinant ATRX antibody.  |
| Produktbeschreibung      | The protein encoded by this gene contains an ATPase/helicase domain, and thus it belongs to the SWI/SNF family of chromatin remodeling proteins. This protein is found to undergo cell cycle-dependent phosphorylation, which regulates its nuclear matrix... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [ATMRX-1R]  |
| UniProt                  | <a href="#">P46100</a>  |
| Reinheit                 | Protein A affinity chromatography   |

|                        |  |
|------------------------|--|
| Formulierung           | 1 mg/ml in 1X PBS, BSA free, sodium azide free   |
| Antibody Type          | Primary Antibody   |
| Application Verdünnung | IHC (FFPE): 1-2ug/ml for 30 min at RT, Prediluted IHC only format: incubate for 30 min at RT (1)   |
| Anwendungsbeschreibung | The concentration stated for each application is a general starting point. Variations in protocols, secondaries and substrates may require the recombinant ATRX antibody to be titered up or down for optimal performance.1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |