

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### PHB Antibody / Prohibitin, Clone: [PBTN-1], Mouse, Monoclonal NSJ-V7598SAF-100UG

|                          |   |
|--------------------------|---|
| Artikelname              | PHB Antibody / Prohibitin, Clone: [PBTN-1], Mouse, Monoclonal   |
| Artikelnummer            | NSJ-V7598SAF-100UG  |
| Hersteller Artikelnummer | V7598SAF-100UG  |
| Alternativnummer         | NSJ-V7598SAF-100UG  |
| Hersteller               | NSJ Bioreagents   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IF, IHC-P   |
| Spezies Reaktivität      | Human   |
| Immunogen                | A recombinant human partial protein corresponding to amino acids 167-261 was used as the immunogen for the PHB antibody.  |
| Produktbeschreibung      | This gene is evolutionarily conserved, and its product is proposed to play a role in human cellular senescence and tumor suppression. Anti-proliferative activity is reported to be localized to the 3 UTR, which is proposed to function as a trans-actin... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [PBTN-1]  |
| UniProt                  | <a href="#">P35232</a>  |
| Reinheit                 | Protein G affinity chromatography   |

|                        |  |
|------------------------|--|
| Formulierung           | 1 mg/ml in 1X PBS, BSA free, sodium azide free   |
| Antibody Type          | Primary Antibody   |
| Application Verdünnung | Immunofluorescence: 1-2ug/ml, Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT, Prediluted IHC only format: incubate for 30 min at RT (1)  |
| Anwendungsbeschreibung | The stated application concentrations are suggested starting points. Titration of the PHB antibody may be required due to differences in protocols and secondary/substrate sensitivity. 1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |