

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD20 Antibody, Clone: [MS4A1/3409], Mouse, Monoclonal NSJ-V7636-100UG

| | |
|--------------------------|---|
| Artikelname | CD20 Antibody, Clone: [MS4A1/3409], Mouse, Monoclonal |
| Artikelnummer | NSJ-V7636-100UG |
| Hersteller Artikelnummer | V7636-100UG |
| Alternativnummer | NSJ-V7636-100UG |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | ELISA, FACS, IF, IHC-P, WB |
| Spezies Reaktivität | Human |
| Immunogen | Amino acids 213-297 from the human protein were used as the immunogen for this CD20 antibody. |
| Produktbeschreibung | Recognizes a protein of 30-33kDa, which is identified as CD20. It is a non-Ig differentiation antigen of B-cells and its expression is restricted to normal and neoplastic B-cells, being absent from all other leukocytes and tissues. CD20 is expressed ... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [MS4A1/3409] |
| UniProt | P11836 |
| Reinheit | Protein G affinity chromatography |

| | |
|------------------------|--|
| Formulierung | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide |
| Antibody Type | Primary Antibody |
| Application Verdünnung | ELISA: 2-4ug/ml (order BSA/azide-free format),Flow cytometry: 1-2ug/10 ⁶ cells,Immunofluorescence: 1-2ug/ml,Western blot: 1-2ug/ml,Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT |
| Anwendungsbeschreibung | The concentration stated for each application is a general starting point. Variations in protocols, secondaries and substrates may require the CD20 antibody to be titered up or down for optimal performance.1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |