

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### CD20 Antibody, Clone: [MS4A1/3410], Mouse, Monoclonal NSJ-V7637-100UG

|                          |   |
|--------------------------|---|
| Artikelname              | CD20 Antibody, Clone: [MS4A1/3410], Mouse, Monoclonal   |
| Artikelnummer            | NSJ-V7637-100UG   |
| Hersteller Artikelnummer | V7637-100UG   |
| Alternativnummer         | NSJ-V7637-100UG   |
| Hersteller               | NSJ Bioreagents   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | ELISA, IHC-P  |
| Spezies Reaktivität      | Human   |
| Immunogen                | Amino acids 213-297 from the human protein were used as the immunogen for this CD20 antibody.   |
| Produktbeschreibung      | Recognizes a protein of 30-33kDa, which is identified as CD20. It is a non-Ig differentiation antigen of B-cells and its expression is restricted to normal and neoplastic B-cells, being absent from all other leukocytes and tissues. CD20 is expressed ... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [MS4A1/3410]  |
| UniProt                  | <a href="#">P11836</a>  |
| Reinheit                 | Protein G affinity chromatography   |

|                        |  |
|------------------------|--|
| Formulierung           | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide   |
| Antibody Type          | Primary Antibody   |
| Application Verdünnung | ELISA (order BSA/sodium azide-free format for coating), Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT   |
| Anwendungsbeschreibung | The concentration stated for each application is a general starting point. Variations in protocols, secondaries and substrates may require the CD20 antibody to be titered up or down for optimal performance.1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |