

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### EpCAM Antibody, Clone: [EGP40/1373], Mouse, Monoclonal NSJ-V7930-20UG

|                          |   |
|--------------------------|---|
| Artikelname              | EpCAM Antibody, Clone: [EGP40/1373], Mouse, Monoclonal  |
| Artikelnummer            | NSJ-V7930-20UG  |
| Hersteller Artikelnummer | V7930-20UG  |
| Alternativnummer         | NSJ-V7930-20UG  |
| Hersteller               | NSJ Bioreagents   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IF, IHC-P, WB   |
| Spezies Reaktivität      | Canine, Feline, Human   |
| Immunogen                | A human partial recombinant protein corresponding to amino acids 100-224 (extracellular domain) was used as the immunogen for the EpCAM antibody.   |
| Produktbeschreibung      | EGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast major... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [EGP40/1373]  |
| UniProt                  | <a href="#">P16422</a>  |
| Reinheit                 | Protein G affinity chromatography   |

|                        |   |
|------------------------|---|
| Formulierung           | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide                |
| Antibody Type          | Primary Antibody  |
| Application Verdünnung | Western blot: 1-2ug/ml,Immunofluorescence: 1-3ug/ml,Immunohistochemistry (FFPE): 1-2ug/ml |
| Anwendungsbeschreibung | Optimal dilution of the EpCAM antibody should be determined by the researcher.            |