

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### CD99 Antibody Cocktail, Clone: [12E7 + MIC2/877], Mouse, Monoclonal NSJ-V7952-20UG

|                          |   |
|--------------------------|---|
| Artikelname              | CD99 Antibody Cocktail, Clone: [12E7 + MIC2/877], Mouse, Monoclonal   |
| Artikelnummer            | NSJ-V7952-20UG  |
| Hersteller Artikelnummer | V7952-20UG  |
| Alternativnummer         | NSJ-V7952-20UG  |
| Hersteller               | NSJ Bioreagents   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | FACS, IF, IHC-P   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Human acute lymphocytic leukemia T-cells (12E7) and recombinant human protein (MIC2/877) were used as the immunogen for the CD99 antibody cocktail.   |
| Produktbeschreibung      | Recognizes a sialoglycoprotein of 27-32kDa, identified as CD99, or MIC2 gene product, or E2 antigen. MIC2 gene is located in the pseudo-autosomal region of the human X and Y chromosome. MIC2 gene encodes two distinct proteins, which are produced by a... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [12E7 + MIC2/877]   |
| UniProt                  | <a href="#">P14209</a>  |

|                        |   |
|------------------------|---|
| Reinheit               | Protein G affinity chromatography   |
| Formulierung           | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide  |
| Antibody Type          | Primary Antibody  |
| Application Verdünnung | Flow cytometry: 1-2ug/10 <sup>6</sup> cells in 0.1ml, Immunofluorescence: 1-2ug/ml, Immunohistochemistry (FFPE): 1-2ug/ml |
| Anwendungsbeschreibung | Optimal dilution of the CD99 antibody should be determined by the researcher.   |