

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Fas Ligand Antibody, Clone: [FASLG/4455], Mouse, Monoclonal NSJ-V8610SAF-100UG

|                          |   |
|--------------------------|---|
| Artikelname              | Fas Ligand Antibody, Clone: [FASLG/4455], Mouse, Monoclonal   |
| Artikelnummer            | NSJ-V8610SAF-100UG  |
| Hersteller Artikelnummer | V8610SAF-100UG  |
| Alternativnummer         | NSJ-V8610SAF-100UG  |
| Hersteller               | NSJ Bioreagents   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC-P   |
| Spezies Reaktivität      | Human   |
| Immunogen                | A portion of amino acids 107-222 from the human protein was used as the immunogen for the Fas Ligand antibody.  |
| Produktbeschreibung      | Cytotoxic T lymphocyte (CTL)-mediated cytotoxicity constitutes an important component of specific effector mechanisms in immunosurveillance against virus-infected or transformed cells. Two mechanisms appear to account for this activity, one of which is... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [FASLG/4455]  |
| UniProt                  | <a href="#">P48023</a>  |
| Reinheit                 | Protein G affinity chromatography   |

|                        |   |
|------------------------|---|
| Formulierung           | 1 mg/ml in 1X PBS, BSA free, sodium azide free                                      |
| Antibody Type          | Primary Antibody  |
| Application Verdünnung | Immunohistochemistry (FFPE): 1-2ug/ml for 30 minutes at RT                          |
| Anwendungsbeschreibung | Optimal dilution of the Fas Ligand antibody should be determined by the researcher. |